
LASER COOLING AND TRAPPING OF NEUTRAL ATOMS(U) NATIONAL BUREAU OF STANDARDS GAITHERSBURG MD ELECTRICITY DIV W D PHILLIPS SEP 84 N00014-84-F-0020 AD-A146 048 1/1 UNCLASSIFIED F/G 20/5 NL END



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

READ INSTRUCTIONS BEFORE COMPLETING FORM

I. REPORT NUMBER

2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER

4. TITLE (and Subtitle)

Laser Cooling and Trapping of Neutral Atoms

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

5. TYPE OF REPORT & PERIOD COVERED INTETIM

1 Oct. 83 - 30 Sept. 84

6. PERFORMING ORG. REPORT NUMBER

7. AUTHOR(a)

William D. Phillips

8. CONTRACT OR GRANT NUMBER(s) N00014-84-F-0020

PERFORMING ORGANIZATION NAME AND ADDRESS National Bureau of Standards

Gaithersburg, MD 20899

10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N

NR 407-007 RR 001-03-03

11. CONTROLLING OFFICE NAME AND ADDRESS

Office of Naval Research Physics Program Office ATTINGTON VA 22217
14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office) 12. REPORT DATE September 1984

13. NUMBER OF PAGES

15. SECURITY CLASS. (of this report)

Unclassified

154. DECLASSIFICATION/ DOWNGRADING SCHEDULE

16. DISTRIBUTION STATEMENT (of this Report)

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, If different from Report)

Approved for public release; distribution unlimited

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

atomic beams, atom traps, frequency standards, high resolution spectroscopy, laser cooling

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)
Using laser cooling techniques, we have produced a sample of stopped atoms with sufficiently large density and small velocity width to be useful for loading neutral atom traps. -

DD 1 JAN 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE 5/N 0102- LF- 014- 6601

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

Annual Summary Report

on

LASER COOLING AND TRAPPING OF NEUTRAL ATOMS

bу


William D. Phillips
National Bureau of Standards
Electricity Division

work supported by
Office of Naval Research
Contract No. N00014-84-F-0020
Task NR 407-007

Date of Report: September 1984

Approved for Public Release; distribution unlimited

Reproduction in whole or in part is permitted for any purpose of the United States Government.



The highest resolution spectroscopy is ultimately limited by processes which relate to atomic velocities such as second order Doppler shift and transit time effects. Laser cooling of ions in traps promises to deal with these problems, but there is a great deal of interest in achieving the same benefits for neutral species. Unfortunately compared to ions, neutrals interact very weakly with external electromagnetic fields, so that it is much more difficult to trap them. Proposed traps using laser fields, electrostatic fields, or magnetic fields, have not yet been realized. Much of the difficulty centers on the very small energy depth of the neutral traps, and the difficulty in cooling atoms contained in such a trap, in contrast to the situation with ion traps. As a result, much of the interest now centers on the deceleration of a free atomic beam, either for direct use in high resolution spectroscopy or as a preparation for loading atoms into a trap.

Deceleration of an atomic beam is accomplished by directing a resonantly tuned laser beam opposite to a beam of neutral atoms. Repeated absorption of the laser light by the atoms, followed by spontaneous re-emission results in deceleration. Use of a magnetic field along with circular polarization of the laser avoids undesired optical pumping of the sodium used in the atomic beam. Causing the magnetic field to vary in space allows the changing Zeeman shift seen by the atoms to compensate the changing Doppler shift as the atoms slow down. In this way, resonance is maintained throughout the deceleration process.

A key problem with the technique described above is that while atoms can be brought to rest, this occurs in a region of strong magnetic field gradient, which makes utilization of the stopped atoms difficult. During the past contract period, we have solved this problem by allowing very slow atoms, created in the gradient field, to drift into a uniform field region after the decelerating laser beam is turned off. Once the atoms are well out of the gradient, the laser is pulsed on for a time sufficient to bring the atoms to rest.

The density of stopped atoms observed exceeds $10^5/\mathrm{cm}^3$, with a velocity width of about 15 m/s. This density and spread should allow the loading of the atoms into a variety of magnetic or optical traps.

Articles published within the past contract period:

- W. D. Phillips, J. V. Prodan, and H. J. Metcalf, "Laser Cooling of Free Neutral Atoms in an Atomic Beam," in Laser Spectroscopy VI, ed. by H. P. Weber and W. Luthy (Springer, Berlin, 1983) p. 162.
- W. D. Phillips and J. V. Prodan, "Cooling Atoms with a Frequency Chirped Laser," in Coherence and Quantum Optics V, ed. by L. Mandel and E. Wolf (Plenum, New York, 1984) p. 15.
- J. V. Prodan, W. D. Phillips and H. J. Metcalf, "Laser Cooling of an Atomic Beam," Conference on Lasers and Electro-optics Technical Digest, Optical Society of America (IEEE cat. No. 83CH1852-3, 1984) p. 34.
- H. J. Metcalf, W. D. Phillips, and J. V. Prodan, "Focussing of Slow Atomic Beams," Bull. Am. Phys. Soc. 29, 795 (1984).

- W. D. Phillips, "Electromagnetic Traps for Neutral Atoms," Bull. Am. Phys. Soc. 29, 815 (1984).
- J. V. Prodan, W. D. Phillips, J. Dalibard and H. J. Metcalf, "Deceleration of Atoms to Zero Velocity," Bull. Am. Phys. Soc. 29, 822 (1984).
- J. V. Prodan, W. D. Phillips, H. J. Metcalf and J. Dalibard, "Stopping Neutral Atoms Using Laser Light," Thirteenth International Quantum Electronics Conference, Technical Digest (IEEE Cat. No. 83-63308) 1984, p. 90.

Publications In Press:

- W. D. Phillips, J. V. Prodan and H. Metcalf "Laser-Cooled Atomic Beams," to appear in Proceedings of the Ninth International Conference on Atomic Physics, Seattle 1984.
- H. J. Metcalf and W. D. Phillips, "Laser Cooling Atomic Beams," to be published in Comments on Atomic and Molecular Physics.
- J. V. Prodan, W. D. Phillips, I. So, H. J. Metcalf, and J. Dalibard, "Stopping Atoms with Laser Light," to be published.

END

FILMED

10-84

DTIC